No.



200400045

THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS; SHAML COME;

Tennessee Adbanced Genetics, Inc.

THE COME, THERE HAS BEEN PRESENTED TO THE

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED DISTINCT VARIETY OF SEXUALLY REPRODUCED, OR TUBER PROPAGATED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF TWENTY YEARS FROM THE DATE OF THIS GRANT, SUBJECT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC REPLENISHMENT OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE RIGHT TO EXCLUDE OTHERS FROM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT, OR IMPORTING IT, OR EXPORTING IT, CONDITIONING IT FOR PROPAGATION, OR STOCKING IT FOR ANY OF THE ABOVE PURPOSE, OR CONDITIONING IT PROPAGATION, OR STOCKING IT FOR ANY OF THE ABOVE PURPOSE, OR USING IT IN PRODUCING A HYBRID OR INT VARIETY THEREFROM, TO THE EXTENT PROVIDED BY THE PLANT VARIETY PROTECTION ACT. IN STATES SEED OF THIS VARIETY (I) SHALL BE SOLD BY VARIETY NAME ONLY AS A CLASS OF CERTIFIED SHALL CONFORM TO THE NUMBER OF GENERATIONS SPECIFIED BY THE OWNER OF THE RIGHTS. (84 STAT. 1542, A ED, 7 U.S.C. 2321 ET SEQ.)

SOYBEAN

'5601T'

In Testimonn Therent, I have hereunto set my hand and caused the seal of the Plant Barrety Protection Office to be affixed at the City of Washington, D.C. this nineteenth day of November, in the year two thousand and four.

Mark

genja

Commissioner Plant Variety Protection Office Agricultural Marketing Service u of Agriculture

NAME (Please print or type)

CAPACITY OR TITLE

ST-470 (02-10-2003) designed by the Plant Variety Protection Office using Word 2000. Replaces former versions of ST-470, which are ob

David N. McKinney

Treasurer

CAPACITY OR TITLE

(See reverse for Instructions and information collection burden statement)

DATE

GENERAL: To be effectively filed with the Plant Variety Protection Office (PVPO), ALL of the following items must application form signed by the owner; (2) completed exhibits A, B, C, E; (3) for a seed reproduced variety at least 2,500 viable untreated seeds, for a hybrid variety at least 2,500 untreated seeds of each line necessary to reproduce the variety, or for tuber reproduced varieties verification that a viable (in the sense that it will reproduce an entire plant) tissue culture will be deposited and maintained in an approved public repository; (4) check drawn on a U.S. bank for \$2,705 (\$320 filling fee and \$2,385 examination fee), payable to "Treasurer of the United States" (See Section 97.6 of the Regulations and Rules of Practice.) Partial applications will be held in the PVPO for not more than 90 days, then returned to the applicant as unfiled. Mail application and other requirements to Plant Variety Protection Office, AMS, USDA, Room 500, NAL Building, 10301 Baltimore Avenue, Beltsville, MD 20705-2351. Retain one copy for your files. All items on the face of the application are self explanatory unless noted below. Corrections on the application form and exhibits must be initialed and dated. DO NOT use masking materials to make corrections. If a certificate is allowed, you will be requested to send a check payable to "Treasurer of the United States" in the amount of \$320 for issuance of the certificate. Certificates will be issued to owner, not licensee or agent.

> Plant Variety Protection Office Telephone: (301) 504-5518 FAX: (301) 504-5291

Homepage: http://www.ams.usda.gov/science/pvpo/pvp.htm

ITEM 18a. Give:

- (1) the genealogy, including public and commercial varieties, lines, or clones used, and the breeding method;
- (2) the details of subsequent stages of selection and multiplication;

(3) evidence of uniformity and stability; and

- (4) the type and frequency of variants during reproduction and multiplication and state how these variants may be identified
- 18b. Give a summary of the variety's distinctness. Clearly state how this application variety may be distinguished from all other varieties in the same crop. If the new variety is most similar to one variety or a group of related varieties:
 - (1) identify these varieties and state all differences objectively;
 - (2) attach statistical data for characters expressed numerically and demonstrate that these are clear differences; and
 - (3) submit, if helpful, seed and plant specimens or photographs (prints) of seed and plant comparisons which clearly indicate distinctness.
- 18c. Exhibit C forms are available from the PVPO Office for most crops; specify crop kind. Fill in Exhibit C (Objective Description of Variety) form as completely
- 18d. Optional additional characteristics and/or photographs. Describe any additional characteristics that cannot be accurately conveyed in Exhibit C. Use comparative varieties as is necessary to reveal more accurately the characteristics that are difficult to describe, such as plant habit, plant color, disease
- 18e. Section 52(5) of the Act requires applicants to furnish a statement of the basis of the applicant's ownership. An Exhibit E form is available from the PVPO.
- 19. If "Yes" is specified (seed of this variety be sold by variety name only, as a class of certified seed), the applicant MAY NOT reverse this affirmative decision after the variety has been sold and so labeled, the decision published, or the certificate issued. However, if "No" has been specified, the applicant may change the choice. (See Regulations and Rules of Practice, Section 97.103).
- See Sections 41, 42, and 43 of the Act and Section 97.5 of the regulations for eligibility requirements.
- 23. See Section 55 of the Act for instructions on claiming the benefit of an earlier filing date.
- 21. CONTINUED FROM FRONT (Please provide a statement as to the limitation and sequence of generations that may be certified.)
- 22. CONTINUED FROM FRONT (Please provide the date of first sale, disposition, transfer, or use for each country and the circumstances, if the variety (including any harvested material) or a hybrid produced from this variety has been sold, disposed of, transferred, or used in the U.S. or other countries.)

January 2003, sold to farmers

23. CONTINUED FROM FRONT (Please give the country, date of filing or issuance, and assigned reference number, if the variety or any component of the variety is protected by intellectual property right (Plant Breeder's Right or Patent).)

NOTES: It is the responsibility of the applicant/owner to keep the PVPO informed of any changes of address or change of ownership or assignment or owner's representative during the life of the application/certificate. There is no charge for filing a change of address. The fee for filing a change of ownership or assignment or any modification of owner's name is specified in Section 97.175 of the regulations. (See Section 101 of the Act, and Sections 97.130, 97.131, 97.175(h) of the Regulations and Rules of Practice.)

To avoid conflict with other variety names in use, the applicant must check the appropriate recognized authority. For example, for agricultural and vegetable crops, contact: Seed Branch, AMS, USDA, Room 213, Building 306, Beltsville Agricultural Research Center--East, Beltsville, MD 20705. Telephone: (301) 504-8089. http://www.ams.usda.gov/lsg/seed/ls-sd.htm

According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a valid OMB control number. The valid OMB control number for this collection of information is (0581-0055). The time required to complete this information collection is estimated to average 1.4 hours per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, sexual orientation, or marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require elternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD). To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call. (202) 720-5964 (voice and TDD). USDA is an equal opportunity provider and employer. S&T-470 (04-01) designed by the Plant Variety Protection Office with WordPerfect 6.0a. Replaces STD-470 (02-99) which is obsolete

EXHIBIT A: ORIGIN AND BREEDING HISTORY (AMENDED JUNE 2004)

5601T is an F₆-derived line from a bi-parental hybridization between 'Hutcheson' [1] and TN89-39. TN89-39 was a selection from the cross TN81-2 x TN80-69. TN81-2 was a selection from the cross 'Lee 74' x 'Mitchell' [2, 3]. TN80-69 was a selection from the cross 'Essex' x J74-40 [8]. J74-40 was a selection from the cross 'Forrest' (2) x D68-18 x PI88788 [5]. D68-18 was a selection from the cross 'Dyer' x 'Bragg' [4, 6]. The F₁ plants from Hutcheson x TN89-39 were grown in Knoxville, TN in 1992. Harvested F₂ seeds were sent to Costa Rica in January 1993 and grown for generation advancement, without selection, via single-pod descent, per plant, to the F₅ stage. The F_{5:6} population was grown in Knoxville, TN in 1994 and agronomically desirable single-plants were harvested. The seeds from these single plants were grown as F_{6.7} plant rows in Knoxville, TN in 1995, and a plant row was identified as visually desirable and bulk harvested. In 1996 seed from that row became designated as the experimental line TN96-58 and was entered in a Preliminary Yield Test grown at two locations (Knoxville and Jackson, TN). TN96-58 was then selected based on superior yield and other desirable agronomic traits. TN96-58 has been evaluated in Tennessee yield tests from 1996 to present. TN96-58 was entered in the Southern Regional Uniform Preliminary Group V Test in 1998 and the Southern Regional Uniform Group V Test in 1999, 2000, and 2001. TN96-58 was designated as the cultivar 5601T upon release in 2001 [7].

The variety 5601T has been reproduced and observed to be uniform and stable, with no variants or off-types, for the past four generations for characteristics listed in this application.

REFERENCES

- 1. Buss, G.R., H.M. Camper, Jr. and C.W. Roane. 1988. Registration of 'Hutcheson' soybean. Crop Sci. 28:1024-1025.
- 2. Caviness, C.E., R.D. Riggs, H.J. Walters. 1975. Registration of 'Lee 74' soybean. Crop Sci. 15:100.
- 3. http://www.ars-grin.gov/cgi-bin/npgs/html/acchtml.pl?1443615. 'Mitchell' soybean is PI 548679.
- 4. Hartwig, E.E. 1968. Registration of 'Dyer' soybean. Crop Sci. 8:402.
- 5. Hartwig, E.E. 1973. Registration of 'Forrest' soybean. Crop Sci. 13:287.
- 6. Hinson, K. and E.E. Hartwig. 1964. Registration of 'Bragg' soybean. Crop Sci. 6:664.
- 7. Pantalone, V.R., F.L. Allen, and D. Landau-Ellis. 2003. Registration of '5601T' soybean. Crop Sci. 43:3 (in press).
- 8. Smith, T.J. and H.M. Camper. 1973. Registration of 'Essex' soybean. Crop Sci. 13:495.

EXHIBIT B: STATEMENT OF DISTINCTNESS (AMENDED MAY 2004)

'5601T' is most similar to 'Hutcheson' except that the **genomic DNA** of '5601T' exhibits a 239 base-pair length band, whereas 'Hutcheson' exhibits a 224 base-pair length band, when amplified via PCR using an M-13 fluorescent type of Satt156 simple sequence repeat primer at the **Satt156 locus of molecular linkage group L**. (See attached Figure P.1).

'5601T' is most similar to 'Hutcheson' except that the **genomic DNA** of '5601T' exhibits a 255 base-pair length band, whereas 'Hutcheson' exhibits a 261 base-pair length band, when amplified via PCR using an M-13 fluorescent type of Satt189 simple sequence repeat primer at the **Satt189 locus of molecular linkage group D1b**. (See attached Figure P.2).

'5601T' is most similar to 'Hutcheson' except that the **genomic DNA** of '5601T' exhibits a 265 base-pair length band, whereas 'Hutcheson' exhibits a 240 base-pair length band, when amplified via PCR using an M-13 fluorescent type of Satt288 simple sequence repeat primer at the Satt288 locus of molecular linkage group G. (See attached Figure P.3).

'5601T' is most similar to 'Hutcheson' except that the **genomic DNA** of '5601T' exhibits a 211 base-pair length band, whereas 'Hutcheson' exhibits a 226 base-pair length band, when amplified via PCR using an M-13 fluorescent type of Satt440 simple sequence repeat primer at the **Satt440 locus of molecular linkage group I.** (See attached Figure P.4).

'5601T' is most similar to 'Hutcheson' except that '5601T' is *significantly taller* in **plant height** than 'Hutcheson'. (See attached Plant Height Tables 1998, 1999, 2000, 2001, 2002, and 1998-2002, note that plant heights are reported in centimeters).

'5601T' is most similar to 'Hutcheson' except that '5601T' is *significantly smaller* in **seed size** than 'Hutcheson'. (See attached Seed Size Tables 1998, 1999, 2000, 2001, 2002, and 1998-2002, note that seed sizes are reported in milligrams per seed).

'5601T' is most similar to 'Hutcheson' except that '5601T' is *significantly higher* in **seed protein concentration** than 'Hutcheson'. (See attached Seed Protein Tables 1998, 1999, 2000, 2001, 2002, and 1998-2002, note that seed protein concentrations are reported in grams of protein per kilogram of seed on a dry-weight basis).

'5601T' is most similar to 'Hutcheson' except that '5601T' is *significantly lower* in **seed oil concentration** than 'Hutcheson'. (See attached Seed Oil Tables 1998, 1999, 2000, 2001, 2002, and 1998-2002, note that seed oil concentrations are reported in grams of oil per kilogram of seed on a dry-weight basis).

'5601T' is most similar to 'Hutcheson' except that '5601T' significantly exceeds 'Hutcheson' in **seed** yield per unit of land area (information previously submitted).

'5601T' is most similar to 'Hutcheson' except that '5601T' is superior to 'Hutcheson' for *tolerance* to Southern root knot nematode *Meloidogyne incognita* (information previously submitted).

'5601T' is most similar to 'Hutcheson' except that '5601T' is *moderately tolerant* to Peanut root knot nematode *Meloidogyne arenaria* while 'Hutcheson' is *highly susceptible*. (information previously submitted).

EXHIBIT B continued: STATEMENT OF DISTINCTNESS

2004 0 004 5

YIELD			Bu	/ ACRE
YEAR	YEAR TEST N LOC		5601T	HUTCHESON
1998	UP VB	9	43.3*	38.7
1999	UT V ¹	24	43.9	43.0
2000	UT V ¹	22	50.1	44.5
2001	UTV	21	51.1	48.5
2002	UTV	18	42.9	40.0
		MEAN	46.7*	43.6

N LOCS = Number of field testing locations

^{* =} significant difference between TN96-58 and Hutcheson at the 0.05 level

MATURITY			days after	HUTCHESON
YEAR	TEST	N LOCS	5601T	HUTCHESON
1998	UP VB	9	+2	0
1999	UTV	24	+2	0
2000	UT V	22	-3	0
2001	UT V	21	-1	0
2002	UTV	18	+0.1	0

HEIGI	HEIGHT			IN	CHES
YE/	\R	TEST	N LOCS	5601T	HUTCHESON
199	8	UP VB	9	29	25
199	9	UTV	24	31	29
200	00	UTV	22	33	30
200)1	UTV	21	32	32
200	2	UTV	18	32	29

PROTEIN			PERCENT PROTEIN	
YEAR	TEST	N LOCS	5601T	HUTCHESON
1998	UP VB	9	42.4	41.1
1999	UTV	24	42.0	40.9
2000	UTV	22	43.3	41.7
2001	UTV	21	41.2	40.1
2002	UTV	18		

SEED SIZE			g / 1	00 seed
YEAR	TEST	NLOCS	5601T	HUTCHESON
1998	UP VB	9	12.6	13.5
1999	UTV	24	13.2	13.9
2000	UTV	. 22	12.3	12.5
2001	UTV	21	13.5	13.8
2002	UT V	18	13.6	13.9

SCN 2			1=RESISTANT	5=SUSCEPTIBLE
YEAR	TEST	LOC	5601T	HUTCHESON
1998	UP VB	TN	5.0	4.1
1999	UTV	TN	5.0	5.0
2000	UTV	TN	5.0	4.9
2001	UTV	TN	4.6	4.7
2002	UTV	TN	4.9	3.8


SCN 3			4.550.55	
			1=RESISTANT	5=SUSCEPTIBLE
YEAR	TEST	LOC	5601T	HUTCHESON
1998	UP VB	TN	4.0	4.7
1999	UTV	TN	5.0	5.0
2000	UTV	TN	4.7	4.7
2001	UTV	TN	5.0	5.0
2002	UTV	TN	3.6	2.1

SCN 14			1=RESISTAN	T 5=SUSCEPTIBLE
YEAR	TEST	LOC	5601T	HUTCHESON
1998	UP VB	TN	4.7	3.3
1999	UTV	TN	4.7	4.3
2000	UTV	TN	4.8	5.0
2001	UTV	TN	4.2	3.8
2002	UT V	TN	4.3	5.0


M. i			1=RESISTANT 5=SUSCEPTIBLE		
YEAR	TEST	LOC	5601T	HUTCHESON	
1998	UP VB	TN	2.3	4.0	
1999	UTV	GA	1.0	5.0	
2000	UTV	GA	1.3	4.5	
2001	UT V	GA	3.8	4.8	
2002	UTV	GA	2.3	5.0	


М. а.		·	1=RESISTANT	5=SUSCEPTIBLE
YEAR	TEST	LOC	5601T	HUTCHESON
1998	UP VB	TN	2.8	3.7
1999	UTV	GA	1.8	5.0
2000	UTV	GA	4.0	4.3
2001	UTV	GA	2.0	3.3
2002	UTV	GA	4.0	4.8


 $^{^{1}}$ = TN96-58 ranked #1 for seed yield among all entries evaluated



7







Form Approved - OMB No. 0581-0055

According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a valid OMB control number. The valid OMB control number for this collection of information is (0581-0055). The time required to complete this information collection is estimated to average 30 minutes per response, including the time for reviewing instructions. searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact the USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call (202) 720-5964 (voice and TDD). USDA is an equal opportunity provider and employer.



U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE SCIENCE AND TECHNOLOGY PLANT VARIETY PROTECTION OFFICE BELTSVILLE, MD 20705

EXHIBIT C (Soybean)

 OBJECTIVE DESCRIPTION OF VARIETY
SOYBEAN (Glycine max (L.) Merr.)

	SUYBEAN (Glyc	ne max (L.) Merr.)	· ·	
NAME OF APPLICANT(S) Tennessee Advanced ADDRESS (Street and No. or R.F.D. No., City, State, and ZIP Code)	Genetics, Inc.		FOR OFFICIAL USE ONLY PVPO NAME OF THE OWNER	0045
2640-C Nolensville Road Nashville, TN 37211			VARIETY NAME 5601T	
			TEMPORARY OR EXPERIMENT DESIGNATION TN96	
PLEASE READ ALL INSTRUCTIONS CAREFULLY: I below.	Place the appropriate nur	nber that describes the var	etal character of this variety	in the boxes
Place a zero in the first box (e.g. 9 9 9 or quantitative	0 9) when	number is either 99 or less	or 9 or less respectively. Data	a for
plant characters should be based on a minimum of 100 pla				trial. Royal
Horticultural Society or any recognized color standard ma Please answer all questions for your variety; lack of respon A. MORPHOLOGY			m used:	
Seed Shape:	`			8
1 = Spherical (L/W, L/T, and T/W ratios < 1.2)		rical-Flattened 1.2; L/T ratio < 1.2)		
3 = Elongate (L/T ratio > 1.2; T/W ratio < 1.2)	4 = Elongate (L/T ratio >	Flattened 1.2;T/W ratio > 1.2)		
Seed Coat Color:			•	,
$1 = Yellow \qquad 2 = Green$	3 = Brown	4 = Black	5 = Other (Please Specify)	·
Seed Coat Luster:				
$\boxed{1} \qquad 1 = \text{Dull} \qquad 2 = \text{Shiny}$		•		
Seed Size:				•
1 3 grams/100 seeds				
Hilum Color:				
$ \begin{array}{ccc} 1 = \text{Buff} & 2 = \text{Yellow} \\ 6 = \text{Black} & 7 = \text{Other (Pleases)} \end{array} $	3 = Brown se Specify)	4 = Gray 5	= Imperfect Black	

Cotyledon Color: mature seed

1 = Yellow

2 = Green

Seed Protein Peroxidase Activity:

1 = Low

2 = High

Hypocotyl Color:

1 = Green

2 = Green with Bronze

('Evans' or 'Davis') Bands below Cotyledon ('Woodworth' or 'Tracy') 3 = Light Purple

below Cotyledons ('Beeson' or 'Pickett 71')

4 = Dark Purple extending to unifoliolate leaves ('Hodgson', 'Coker', or 'Hampton 266A')

Leaf Shape:

1 = Lanceolate

2 = Oval

3 = Ovate

4 = Other (Please Specify)

Flower Color:

1 = White

2 = Purple

3 = White with a Purple Throat

Pod Color:

1 = Tan

2 = Brown

3 = Black

Pubescence Color:

1 = Grav

2 = Brown (Tawny)

3 = Light Tawny

Plant Habit:

1 = Determinate

2 = Semi - Determinate

3 = Indeterminate

4 = Intermediate

Maturity Group:

1 = 0006 = III11 = VIII 2 = 00

7 = IV

12 = IX

3 = 0

8 = V13 = X 4 = I

9 = VI

14 = XI

10 = VII15 = XII

5 = II

Maturity Subgroup:

6 | Please enter a value from 0 - 9

B. DISEASE REACTIONS

0 = Not Tested

1 = Susceptible

2 = Resistant

3 = Tolerant

Bacterial

Bacterial Pustule (Xanthomonas campestris pv. glycines (Nakano) Dye)

Bacterial Blight (Pseudomonas syringae pv. glycinea (Coerper) Young, Dye, & Wilkie)

Wildfire Blight (Pseudomonas syringae pv. tabaci (Wolf & Foster) Young, Dye, & Wilkie)

B. 3	DISEASE REACTIONS (Continu	$\theta = \text{Not Tested}$	1 = Susceptible	2 = Resistant 2004er 0t 004 5
Fung	gal		м .	
C	Brown Spot (Septoria glycines	Hemmi)		
	Frogeye Leaf Spot (Cercospor	a sojina Hara)		
	race 1	race 2	race 3	race 4
	race 5	race 6		(Please Specify)
0	Target Spot (Corynespora cass	siicola (Berk. & Curt.) Wei)	in Mi	cally occuring population ilan Tennessee
0	Downey Mildew (Peronospora	trifoliorum var. manchurica	τ (Naum.) Syd. ex G	äum)
0	Powdery Mildew (Microsphae)	ra diffusa Cke. & Pk.)		
0	Brown Stem Rot (Phialophora	gregata (Allington & Chaml	perlain) W. Gams.)	. •
2	Stem Canker (Diaporthe phase	olorum (Cke. & Ell.) Sacc. va	ar. <i>caulivora</i> Athow	& Caldwell)
0	Pod and Stem Blight (<i>Diaporth</i>	e phaseolorum (Cke. & Ell.)	Sacc. var. sojae (Le	hman) Wehm.)
0	Purple Seed Stain (Cercospora	kikuchii (T. Matsu. & Tomo	yasu) Gardener)	
0	Rhizoctonia Root Rot (Rhizocto	onia solani Kühn)		
hytop	ohthora Root Rot (<i>Phytophthora</i> i	negasperma Drechs. f. sp. gly	vcinea (Kuan & Erw	rin))
0	race 1 O race 8	O race 15	race 22	
0	race 2 O race 9	0 race 16	race 23	
0	race 3 race 10	race 17	race 24	
0	race 4 race 11	o race 18	race 25	
0	race 5 race 12	race 19	race 26	
0	race 6 0 race 13	0 race 20 0	Other (Please Si	pecify)
0	race 7 race 14	race 21		
0	Bud Blight (Tobacco Ringspot			
0	Yellow Mosaic (Bean Yellow M	osaic Virus)		

В. 1	DISEASE REACTIONS (Continued)	0 = Not Tested	1 = Susceptible	2 = Resistant	0.0 401e AnO	04
. 0	Cowpea Mosaic (Cowpea Chlorotic	c Virus)	•			٠
0	Pod Mottle (Bean Pod Mottle Virus	s)	÷		•	
2	Seed Mottle (Soybean Mosaic Virus	s)				
Nema	tode		·			
Soybe	ean Cyst Nematode (Heterodera glycine	es Ichinohe)	•			
0	race 1 0 race 4	0 race 9	•	•.		
1	race 2 O race 5	1 race 1	14			
1	race 3 orace 6	Other	(Please Specify)	•		
0	Lance Nematode (Hoplolaimus colu	mbus Sher)			÷	
3	Southern Root Knot Nematode (Med	loidogyne incognita (I	Kofoid & White) Chit	wood)		
0	Northern Root Knot Nematode (Med	loidogyne hapla Chity	wood)			
3	Peanut Root Knot Nematode (Meloid	dogyne arenaria (Nea	I) Chitwood)			
0	Reniform Nematode (Rotylenchus re	niformus Linwood &	Olivera)		· .	
0	Javanese Nematode (Meloidogyne ja	vanica (Treub) Chitw	vood)			
0	Other Nematode (Please Specify)	· .				
C. PH	YSIOLOGICAL RESPONSES	0 = Not Tested	1 = Susceptible	2 = Resistant	3 = Tolerant	
0	Iron Chlorosis on Calcareous Soil					
0	Phosphorus	Other	(Please Specify)		· · · · · · · · · · · · · · · · · · ·	
0	Boron		•			
0	Aluminum					
0	Salt					•
0	Drought					

S&T-470-2 (2-90) decianed by the Plant Variation Dec

D. INSECT REACTIONS	0 = Not Tested	1 = Susceptible	2 = Resistant	UUU43 3 = Tolerant
O Mexican Bean Beetle (Epilachna va	rivestis Mulsant)	*		
O Potato Leaf Hopper (Empoasca fab	ae (Harris))			
O Other (Please Specify)				
E. HERBICIDE REACTIONS	0 = Not Tested	1 = Susceptible	2 = Resistant	
O Metribuzin				
O Bentazone				
O Sulfonylurea				
1 Glyphosate				
O Glufosinate			,	
O Pendimethalin				
O Other (Please Specify)		•		
F. TRANSGENIC COMPOSITION				
Has the development of the subject variety in or, the removal of genetic material from the a f yes, please complete the following informat	application variety?	· '		er than a soybean,
. Please state the vector's name:				
. Please state the vector components:			· (
. Please describe the genetic material succe	essfully transferred in	to the subject variety	:	
. Please describe the insertion protocol:				
A literature citation(s) explaining the fou the "Transgenic Composition" portion of	r information request this form.	s above may be an ac	ceptable alternative	to completion of

Please describe any biochemical information here, which you believe will be helpful in further describing the subject variety (e.g. Simple Sequence Repeats, Restriction Fragment Length Polymorphisms, Isozymic Characterization). Use additional pages if necessary.

G. BIOCHEMICAL MARKERS

H. COMMENTS

Plant Height Table 1998. Mean plant height, per location, in the 1998 USDA-ARS Southern Uniform Test.

200400045

	Plant Height (cm)		
1998 Location	'5601T'	'Hutcheson	
BIXBY,OK98	76.2	59.7	
JACKSON,TN98	86.4	74.9	
KEISER,AR98	74.9	60.3	
PITTSBURG,KS98	105.4	94.6	
PLYMOUTH,NC98	87.6	68.6	
PORTAGEVILLE,MO(A)98	63.5	67.3	
QUEENSTOWN,MD98	88.9	71.1	
STONEVILLE,MS98	33.0	27.9	
ULLIN,IL98	59.7	56.5	
WARSAW,VA98	63.5	59.0	
1998 Mean	73.9	64.0	

Statistical analysis via ANOVA, '5601T' differs significantly from 'Hutcheson'

F-value = 19.2, P-value = 0.0018

 $R^2 = 96.6\%$, CV = 8.4%Uniform test plots typically consist of four-row plots, 20 feet in length, replicated 3-times per location.

Plant Height Table 1999. Mean plant height, per location, in the 1999 USDA-ARS Southern Uniform Test.

·	Plant Height (cm)		
1999 Location	'5601T'	'Hutcheson	
BELLE MINA,AL99	69.4	78.7	
BIXBY,OK99	61.0	55.9	
BOSSIER CITY,LA99	80.4	73.7	
GEORGETOWN,DE99	114.3	81.3	
Knoxville,TN99	80.4	74.5	
MCCUNE,KS99	51.6	50.0	
PINE TREE,AR99	59.3	61.8	
PLYMOUTH,NC99	87.2	77.9	
PORTAGEVILLE,MO(A)99	76.2	76.2	
PORTAGEVILLE,MO(B)99	71.10	68.6	
PRINCETON,KY99	109.20	92.7	
PROSPER,TX99	33.0	40.60	
QUEENSTOWN,MD99	83.80	73.7	
SPRINGFIELD,TN99	71.1	66.9	
STONEVILLE,MS99	71.1	66.0	
STUTTGART,AR99	94.0	88.9	
SUFFOLK,VA99	88.0	81,3	
ULLIN,IL99	86.40	93.1	
WARSAW,VA99	61.0	63.5	
1999 Mean	76.2	71.8	
Statistical analysis via ANOVA, '5601T' differs			
significantly from 'Hutcheson'			

significantly from 'Hutcheson'

F-value = 4.2, P-value =0.05

R² = 93.5%, CV = 10.6%

Uniform test plots typically consist of four-row plots, 20 feet in length, replicated 3-times per location.

Plant Height Table 2000. Mean plant height, per location, in the 2000 USDA-ARS Southern Uniform Test.

Plant Height (cm)		
'5601T'	'Hutcheson	
80.4	69.8	
68.6	58.4	
72.0	66.0	
111.8	99.1	
88.0	87.2	
78.7	63.5	
99.1	82.1	
96.5	86.4	
80.4	74.5	
50.8	50.8	
114.3	104.1	
94.0	83.80	
76.2	63.5	
76.2	66.0	
45.7	71.1	
78.7	71.1	
52.5	50.0	
98.2	96.5	
89.7	83.0	
81.7	75.1	
	*5601T' 80.4 68.6 72.0 111.8 88.0 78.7 99.1 96.5 80.4 50.8 114.3 94.0 76.2 76.2 45.7 78.7 52.5 98.2 89.7	

Statistical analysis via ANOVA, '5601T' differs significantly from 'Hutcheson'
F-value = 12.67, P-value = 0.0022 $R^2 = 98.0\%$, CV = 6.0%Uniform test plots typically consist of four-row plots,

20 feet in length, replicated 3-times per location.

Plant Height Table 2001. Mean plant height, per location, in the 2001 USDA-ARS Southern Uniform Test.

	Plant Height (cm)			
2001 Location	'5601T'	'Hutcheson		
BELLE MINA,AL01	76.2	83.0		
BOSSIER CITY,LA01	73.7	62.6		
GEORGETOWN,DE01	88.9	89.7		
KEISER,AR01	96.5	101.6		
Knoxville,TN01	77.0	76.2		
MCCUNE,KS01	75.4	61.8		
ORANGE,VA01	110.1	111.8		
PINE TREE,AR01	104.1	96.5		
PITTSBURG,KS01	72.0	64.3		
PLYMOUTH,NC01	81.3	71.1		
PORTAGEVILLE,MO(A)01	73.7	83.8		
PORTAGEVILLE,MO(B)01	92.3	88.0		
PRINCETON,KY01	107.5	99.9		
PROSPER,TX01	58.4	48.3		
QUEENSTOWN,MD01	86.4	76.2		
SPRINGFIELD,TN01	116.0	110.1		
STARKVILLE,MS01	63.5	61.0		
STONEVILLE,MS01	81.3	101.6		
STUTTGART,AR01	41.9	52.1		
SUFFOLK,VA01	67.7	71.1		
ULLIN,IL01	72.8	88.9		
WARSAW,VA01	86.4	80.4		
2001 Mean	82.0	80.9		
Statistical analysis via ANOVA,				
No significant difference in 200)1			

Uniform test plots typically consist of four-row plots, 20 feet in length, replicated 3-times per location.

Plant Height Table 2002. Mean plant height, per location, in the 2002 USDA-ARS Southern Uniform Test.

	Plant Height (cm)		
2002 Location	'5601T'	'Hutcheson	
BELLE MINA,AL02	66.0	63.5	
BIXBY,OK02	76.2	66.0	
BOSSIER CITY,LA02	62.6	55.0	
GEORGETOWN,DE02	113.4	109.2	
KEISER,AR02	100.8	88.0	
Knoxville,TN02	94.8	74.5	
MARIANNA,AR02	84.7	87.2	
MCCUNE,KS02	90.6	79.6	
ORANGE,VA02	80.4	78.7	
PITTSBURG,KS02	66.9	57.6	
PORTAGEVILLE,MO(A)02	81.3	71.1	
PORTAGEVILLE,MO(B)02	71.1	61.0	
PRINCETON,KY02	104.1	94.0	
PROSPER,TX02	39.8	35.6	
QUEENSTOWN,MD02	90.6	77.9	
STONEVILLE,MS02	61.0	61.0	
SUFFOLK,VA02	75.4	70.3	
ULLIN,IL02	83.8	88.9	
WARSAW,VA02	75.4	64.3	
2002 Mean	79.9	72.8	
Statistical analysis via ANOVA, '5601T' differs			
significantly from 'Hutcheson'			
T 1 10.70 D 1 0.0002			

F-value = 19.72, P-value = 0.0003 R² = 95.2%, CV = 9.5% Uniform test plots typically consist of four-row plots, 20 feet in length, replicated 3-times per location.

	Plant Height (cm)	
Year	'5601T'	'Hutcheson
1998	73.9	64.0
1999	76.2	71.8
2000	81.7	75.1
2001	82.0	80.9
2002	79.9	72.8
Mean 1998-2002	78.7	72.9
Statistical analysis via ANOVA '5601T' differs		

Statistical analysis via ANOVA, significantly from 'Hutcheson'

F-value = 30.9, P-value =0.0001

R² = 96.1%, CV = 8.4%

Uniform test plots typically consist of four-row plots, 20 feet in length, Replicated 3-times per environment.

Seed Size Table 1998. Mean Seed Size, per location, in the 1998 USDA-ARS Southern Uniform Test.

	Seed Size (mg/seed)		
1998 Location	'5601T'	'Hutcheson	
BIXBY,OK98	145.0	157.0	
JACKSON,TN98	131.0	133.2	
PITTSBURG,KS98	96.0	112.0	
PLYMOUTH,NC98	140.0	154.0	
PORTAGEVILLE,MO(A)98	105.0	121.5	
QUEENSTOWN,MD98	141.5	161.2	
STONEVILLE,MS98	107.0	122.0	
ULLIN,IL98	99.0	108.2	
WARSAW98	135.5	133.8	
1998 Mean	122.2	133.7	

Statistical analysis via ANOVA, '5601T' differs significantly from 'Hutcheson'

F-value = 17.1, P-value = 0.0033 R² = 94.5%, CV = 5.3%

Uniform test plots typically consist of four-row plots, 20 feet in length, replicated 3-times per location.

Seed Size Table 1999. Mean Seed Size, per location, in the 1999 USDA-ARS Southern Uniform Test.

	Seed Size (mg/seed)		
1999 Location	'5601T'	'Hutcheson	
BELLE MINA,AL99	98.0	117.0	
BIXBY,OK99	146.0	167.0	
Knoxville,TN99	118.0	138.0	
MCCUNE,KS99	136.0	134.0	
PINE TREE,AR99	166.0	162.3	
PLYMOUTH,NC99	114.0	125.0	
PORTAGEVILLE,MO(A)99	132.0	133.0	
PORTAGEVILLE,MO(B)99	121.0	134.0	
PRINCETON,KY99	140.0	125.00	
PROSPER,TX99	96.0	109.0	
QUEENSTOWN,MD99	118.3	138.0	
SPRINGFIELD,TN99	112.0	132.0	
STONEVILLE,MS99	143.0	152.0	
SUFFOLK,VA99	98.0	109.3	
ULLIN,IL99	102.0	108.3	
WARSAW,VA99	147.0	156.0	
1999 Mean	124.2	133.7	

Statistical analysis via ANOVA, '5601T' differs significantly from 'Hutcheson'

F-value = 12.24, P-value = 0.0032 $R^2 = 97.7$ %, CV = 6.4% Uniform test plots typically consist of four-row plots, 20 feet in length, replicated 3-times per location.

Seed Size Table 2000. Mean Seed Size, per location, in the 2000 USDA-ARS Southern Uniform Test.

	Seed Size (mg/seed)			
2000 Location	'5601T'	'Hutcheson		
BELLE MINA,AL00	106.0	123.0		
CALHOUN,GA00	140.0	140.0		
Knoxville,TN00	133.0	143.0		
MCCUNE,KS00	109.0	118.0		
ORANGE,VA00	150.3	158.3		
PITTSBURG,KS00	99.0	112.0		
PORTAGEVILLE,MO(A)00	111.0	134.0		
PORTAGEVILLE,MO(B)00	122.0	135.0		
PRINCETON,KY00	135.0	145.0		
QUEENSTOWN,MD00	145.0	161.0		
SPRINGFIELD,TN00	123.0	130.0		
STONEVILLE,MS00	122.0	112.0		
SUFFOLK,VA00	139.0	163.7		
ULLIN,IL00	131.7	127.3		
WARSAW,VA00	142.7	140.0		
2000 Mean	127.2	136.2		
Statistical analysis via ANOVA, '5601T' differs				
significantly from 'Hutcheson'				
F-value = 8.36, P -value = 0.0119				
$P^2 = 06.80\%$ CV = 5.20%				

 $R^2 = 96.8\%$, CV = 5.2%

Uniform test plots typically consist of four-row plots,

20 feet in length, replicated 3-times per location.

Seed Size Table 2001. Mean Seed Size, per location, in the 2001 USDA-ARS Southern Uniform Test.

	Seed Size (mg/seed)	
2001 Location	'5601T'	'Hutcheson
BELLE MINA,AL01	182.3	168.3
Knoxville,TN01	118.0	130.0
MCCUNE,KS01	139.0	151.0
ORANGE,VA01	116.0	128.3
PINE TREE,AR01	150.3	153.3
PITTSBURG,KS01	136.0	142.0
PLYMOUTH,NC01	143.0	154.0
PORTAGEVILLE,MO(A)01	128.0	142.0
PORTAGEVILLE,MO(B)01	140.0	145.0
PRINCETON,KY01	127.0	118.0
QUEENSTOWN,MD01	124.0	123.7
STONEVILLE,MS01	109.0	109.0
SUFFOLK,VA01	125.3	124.3
ULLIN,IL01	127.0	125.0
WARSAW,VA01	139.7	148.3
2001 Mean	133.6	137.5
Statistical analysis via ANOVA,		
No significant difference in 20		

Uniform test plots typically consist of four-row plots, 20 feet in length, replicated 3-times per location.

Seed Size Table 2002. Mean Seed Size, per location, in the 2002 USDA-ARS Southern Uniform Test.

	Seed Size (mg/seed)	
2002 Location	'5601T'	'Hutcheson
BIXBY,OK02	163.0	137.0
Knoxville,TN02	123.0	143.0
MCCUNE,KS02	125.0	132.0
ORANGE, VA02	183.3	186.7
PITTSBURG,KS02	116.0	111.0
PLYMOUTH,NC02	157.0	174.0
PORTAGEVILLE,MO(A)02	123.0	121.0
PORTAGEVILLE,MO(B)02	111.0	107.0
PRINCETON,KY02	108.1	111.2
PROSPER,TX02	119.0	133.0
SUFFOLK,VA02	165.7	171.0
ULLIN,IL02	143.7	148.3
WARSAW,VA02	155.7	170.7
2002 Mean	138.0	142.0
Statistical analysis via ANOVA,		
No significant difference in 20	01	

Uniform test plots typically consist of four-row plots, 20 feet in length, replicated 3-times per location.

Seed Size Table 1998-2002. Five-year mean Seed Size in the USDA-ARS Southern Uniform Tests

	Seed Siz	Seed Size (mg/seed)	
Year	'5601T'	'Hutcheson	
1998	122.2	133.7	
1999	124.2	133.7	
2000	127.2	136.2	
2001	133.6	137.5	
2002	138.0	142.0	
Mean 1998-2002	129.0	136.6	
G: .: 1 1 1 1	1310111 (#60	4 500 74 60	

Statistical analysis via ANOVA, '5601T' differs

significantly from 'Hutcheson'

F-value = 31.2, P-value = 0.0001

R² = 96.8%, CV = 5.9%

Uniform test plots typically consist of four-row plots, 20 feet in length, Replicated 3-times per environment.

Seed Protein Concentration Table 1998. Mean Seed Protein Concentration, per location in the 1998 USDA ARS Seed Protein Concentration.

per location, in the 1998 USDA-ARS Southern Uniform Test.

	Seed Protein Concentration (g/kg dry weight)	
1998 Location	'5601T'	'Hutcheson
JACKSON,TN98	435	424
PLYMOUTH,NC98	423	398
PORTAGEVILLE,MO(A)98	440	428
QUEENSTOWN,MD98	419	389
STONEVILLE,MS98	406	392
ULLIN,IL98	435	426
WARSAW,VA98	412	416
1998 Mean	424	411

Statistical analysis via ANOVA, '5601T' differs significantly from 'Hutcheson'

F-value = 10.71, P-value = 0.0170

 $R^2 = 96.9\%$, CV = 1.2%Uniform test plots typically consist of four-row plots, 20 feet in length, replicated 3-times per location.

Seed Protein Concentration Table 1999. Mean Seed Protein Concentration, per location, in the 1999 USDA-ARS Southern Uniform Test.

	Seed Protein Concentration (g/kg dry weight)	
1999 Location	'5601T'	'Hutcheson
BELLE MINA,AL99	454	442
Knoxville,TN99	442	426
MCCUNE,KS99	364	381
PINE TREE,AR99	432	408
PLYMOUTH,NC99	421	411
PORTAGEVILLE,MO(A)99	419	410
PRINCETON,KY99	449	403
PROSPER,TX99	438	450
QUEENSTOWN,MD99	434	429
STONEVILLE,MS99	431	408
SUFFOLK,VA99	429	433
ULLIN,IL99	469	468
WARSAW,VA99	383	376
1999 Mean	428	419

Statistical analysis via ANOVA, '5601T' differs from

'Hutcheson' but at a 0.10 probability level.

F-value = 4.13, P-value = 0.0649

 $R^2 = 91.1\%$, CV = 2.7%

Uniform test plots typically consist of four-row plots, 20 feet in length, replicated 3-times per location.

Seed Protein Concentration Table 2000. Mean Seed Protein Concentration, per location, in the 2000 USDA-ARS Southern Uniform Test.

	Seed Protein Concentration (g/kg dry	
	weight)	
2000 Location	'5601T'	'Hutcheson
BELLE MINA,AL00	414	405
Knoxville,TN00	420	407
ORANGE,VA00	421	405
PITTSBURG,KS00	459	435
PORTAGEVILLE,MO(A)00	430	415
PRINCETON,KY00	442	426
QUEENSTOWN,MD00	444	426
SPRINGFIELD,TN00	415	403
STONEVILLE,MS00	457	418
SUFFOLK,VA00	413	405
ULLIN,IL00	440	423
WARSAW,VA00	436	412
2000 Mean	433	415

Statistical analysis via ANOVA, '5601T' differs significantly from 'Hutcheson'

F-value = 52.94, P-value = 0.0001

 $R^2 = 94.3\%$, CV = 1.4%

Uniform test plots typically consist of four-row plots, 20 feet in length, replicated 3-times per location.

Seed Protein Concentration Table 2001. Mean Seed Protein Concentration,

per location in the 2001 HSDA ARS Seed.

per location, in the 2001 USDA-ARS Southern Uniform Test.

	Seed Protein Concentration (g/kg dry weight)	
2001 Location	'5601T'	'Hutcheson
Knoxville,TN01	429	411
MCCUNE,KS01	396	400
ORANGE,VA01	404	393
PLYMOUTH,NC01	417	400
PORTAGEVILLE,MO(A)01	427	409
PRINCETON,KY01	404	386
PROSPER,TX01	408	418
QUEENSTOWN,MD01	409	409
STONEVILLE,MS01	415	398
ULLIN,IL01	421	408
WARSAW,VA01	393	387
2001 Mean	411	402

Statistical analysis via ANOVA, '5601T' differs significantly from 'Hutcheson'

F-value = 9.75, P-value = 0.0108 $R^2 = 82.8\%$, CV = 1.7%Uniform test plots typically consist of four-row plots, 20 feet in length, replicated 3-times per location.

Seed Protein Concentration Table 2002. Mean Seed Protein Concentration,

per location in the 2002 HSDA ARS Seed Protein Concentration,

per location, in the 2002 USDA-ARS Southern Uniform Test.

	Seed Protein Concentration (g/kg dry weight)	
2002 Location	'5601T'	'Hutcheson
BIXBY,OK02	419	420
Knoxville,TN02	419	394
MCCUNE,KS02	388	397
ORANGE,VA02	382	377
PLYMOUTH,NC02	411	403
PORTAGEVILLE,MO(A)02	402	381
PRINCETON,KY02	435	425
PROSPER,TX02	426	418
STONEVILLE,MS02	401	391
ULLIN,IL02	406	387
WARSAW,VA02	429	430
2002 Mean	411	402
Statistical analysis via ANOVA, '5601T' differs		

significantly from 'Hutcheson'

F-value = 7.87, P-value = 0.0186 $R^2 = 92.1\%$, CV = 1.8%Uniform test plots typically consist of four-row plots, 20 feet in length, replicated 3-times per location.

	Seed Protein Concentration (g/kg dry weight)	
Year	'5601T'	'Hutcheson
1998	424	411
1999	428	419
2000	433	415
2001	411	402
2002	411	402
Mean 1998-2002	421	410

Statistical analysis via ANOVA, '5601T' differs significantly from 'Hutcheson'

F-value = 51.53, P-value = 0.0001

 $R^2 = 99.6\%$, CV = 1.2%Uniform test plots typically consist of four-row plots, 20 feet in length, Replicated 3-times per environment.

Seed Oil Concentration Table 1998. Mean Seed Oil Concentration, per location, in the 1998 USDA-ARS Southern Uniform Test.

	Seed Oil Concentration (g/kg dry weight)	
1998 Location	'5601T'	'Hutcheson
JACKSON,TN98	198	207
PLYMOUTH,NC98	196	210
PORTAGEVILLE,MO(A)98	189	201
QUEENSTOWN,MD98	195	214
STONEVILLE,MS98	196	219
ULLIN,IL98	190	203
WARSAW,VA98	209	212
1998 Mean	196	209

Statistical analysis via ANOVA, '5601T' differs significantly from 'Hutcheson'

F-value = 27.55, P-value = 0.0019

 $R^2 = 91.9\%$, CV = 2.1%

Uniform test plots typically consist of four-row plots, 20 feet in length, replicated 3-times per location.

Seed Oil Concentration Table 1999. Mean Seed Oil Concentration, per location, in the 1999 USDA-ARS Southern Uniform Test.

20040004	
----------	--

	Seed Oil Concentration	
	(g/kg dry weight)	
1999 Location	'5601T'	'Hutcheson
BELLE MINA,AL99	181	197
Knoxville,TN99	197	212
MCCUNE,KS99	187	193
PINE TREE,AR99	196	212
PLYMOUTH,NC99	187	199
PORTAGEVILLE,MO(A)99	194	204
PRINCETON,KY99	175	197
PROSPER,TX99	183	189
QUEENSTOWN,MD99	187 .	192
STONEVILLE,MS99	192	211
SUFFOLK,VA99	180	187
ULLIN,IL99	173	183
WARSAW,VA99	202	201
1999 Mean	187	198
Statistical analysis via ANOVA	, '5601T' di	iffers
significantly from 'Hutcheson'		

Uniform test plots typically consist of four-row plots, 20 feet in length, replicated 3-times per location.

F-value = 37.75, P-value = 0.0001

 $R^2 = 91.1 \%$, CV = 2.4

Seed Oil Concentration Table 2000. Mean Seed Oil Concentration, per location, in the 2000 USDA-ARS Southern Uniform Test. 200400045

	Seed Oil Concentration (g/kg dry weight)	
2000 Location	'5601T'	'Hutcheson
BELLE MINA,AL00	197	207
Knoxville,TN00	202	217
ORANGE,VA00	197	199
PITTSBURG,KS00	185	197
PORTAGEVILLE,MO(A)00	194	200
PRINCETON,KY00	193	204
QUEENSTOWN,MD00	186	196
SPRINGFIELD,TN00	191	212
STONEVILLE,MS00	194	207
SUFFOLK,VA00	200	195
ULLIN,IL00	194	· 205
WARSAW,VA00	193	208
2000 Mean	194	204

Statistical analysis via ANOVA, '5601T' differs significantly from 'Hutcheson'

F-value = 27.34, P-value = 0.0003

 $R^2 = 82.6\%$, CV = 2.4%Uniform test plots typically consist of four-row plots, 20 feet in length, replicated 3-times per location.

Seed Oil Concentration Table 2001. Mean Seed Oil Concentration, per location, in the 2001 USDA-ARS Southern Uniform Test.

	Seed Oil Concentration (g/kg dry weight)		
2001 Location	'5601T'	'Hutcheson	
Knoxville,TN01	205	217	
MCCUNE,KS01	216	212	
ORANGE,VA01	202	211	
PLYMOUTH,NC01	201	212	
PORTAGEVILLE,MO(A)01	210	214	
PRINCETON,KY01	207	220	
PROSPER,TX01	218	209	
QUEENSTOWN,MD01	193	201	
STONEVILLE,MS01	209	220	
ULLIN,IL01	209	216	
WARSAW,VA01	198	215	
2001 Mean	206	213	
Statistical analysis via ANOVA, '5601T' differs significantly from 'Hutcheson'			

F-value = 9.72, P-value = 0.0109 $R^2 = 74.2\%$, CV = 2.6%

Uniform test plots typically consist of four-row plots, 20 feet in length, replicated 3-times per location.

2004000

Seed Oil Concentration Table 2002. Mean Seed Oil Concentration, per location, in the 2002 USDA-ARS Southern Uniform Test.

	Seed Oil Concentration (g/kg dry weight)	
2002 Location	'5601T'	'Hutcheson
BIXBY,OK02	183	191
Knoxville,TN02	207	217
MCCUNE,KS02	208	203
ORANGE,VA02	189	194
PLYMOUTH,NC02	191	199
PORTAGEVILLE,MO(A)02	192	201
PRINCETON,KY02	199	204
PROSPER,TX02	193	203
STONEVILLE,MS02	186	214
ULLIN,IL02	197	211
WARSAW,VA02	189	189
2002 Mean	194	202

Statistical analysis via ANOVA, '5601T' differs significantly from 'Hutcheson'

F-value = 11.08, P-value = 0.0076R² = 81.2%, CV = 3.0%

Uniform test plots typically consist of four-row plots, 20 feet in length, replicated 3-times per location.

Seed Oil Concentration Table 1998-2002. Five-year mean

Seed Oil Concentration	in the USDA-ARS Southern Uniform Tests

	Seed Oil Concentration (g/kg dry weight)	
Year	'5601T'	'Hutcheson
1998	196	209
1999	187	198
2000	194	204
2001	206	213
2002	194	202
Mean 1998-2002	196	205

Statistical analysis via ANOVA, '5601T' differs

significantly from 'Hutcheson'

F-value = 99.13, P-value =0.0001 $R^2 = 99.0\%$, CV = 2.2%Uniform test plots typically consist of four-row plots, 20 feet in length, Replicated 3-times per environment.

REPRODUCE LOCALLY, Include form number and edition date on a	di roproduce.	
U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE EXHIBIT E	Application is required in order to determine if a plant variety protection certificate is to be issued (7/1/S C 2/131). The information	
STATEMENT OF THE BASIS OF OWNERSHIP 1. NAME OF APPLICANT(S)	and the certificate is iss	sued (7 U.S.C. 2426).
	2. TEMPORARY DESIGNATION OR EXPERIMENTAL NUMBER	3. VARIETY NAME
Tennessee Advanced Genetics, Inc.	TN96-58	5601T
4. ADDRESS (Street and No., or R.F.D. No., City, State, and ZIP, and Country)	5. TELEPHONE (Include area code)	6. FAX (Include area code)
2640-C Nolensville Road Nashville, TN 37211	615-242-0467	615-248-3461
	7. PVPO NUMBER 2004 0	0045
8. Does the applicant own all rights to the variety? Mark an "X" in the	appropriate block. If no, please exp	lain YES NO
		X YES NO
9 is the applicant finding in		·
9. Is the applicant (individual or company) a U.S. National or a U.S. ba	ased company? If no, give name of c	ountry YES NO
10. Is the applicant the original owner? YES X NO	If no, please answer one of the fol	louis a
. — —		
a. If the original rights to variety were owned by individual(s), is (ar	re) the original owner(s) a U.S. Nationa	al(s)?
YES NO	If no, give name of country	
b. If the original rights to vortate was		
 b. If the original rights to variety were owned by a company(ies), is 	s (are) the original owner(s) a U.S. bas	ed company?
X YES NO	If no, give name of country	
11. Additional explanation on ownership (If needed, use the reverse for	eytra enacol:	
•	· ·	
Acquired from the University of Tennes	see, knoxville, in 3/990	6
		•
PLEASE NOTE:		
Plant variety protection can only be afforded to the owners (not licensees	s) who meet the following criteria:	•
 If the rights to the variety are owned by the original breeder, that personational of a country which affords similar protection to nationals of the 	The second of the second seconds	•
If the rights to the variety are owned by the company which employed nationals of a UPOV member country, or owned by nationals of a cour genus and species.		
3. If the applicant is an owner who is not the original owner, both the original owner, both the original owner.		
The original breeder/owner may be the individual or company who directed Act for definitions.	ed the final breeding. See Section 41(a)(2) of the Plant Variety Protection
According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, and a control number. The valid OMB control number for this information collection is 0581-0055. The tresponse, including the time for reviewing the instructions, searching existing data sources, gather The U.S. Department of Agriculture (USDA) prohibits discontrolled in the	find and maintaining the data and midreton conection	is estimated to average 6 minutes per
The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs on the basis and marital or familial status. (Not all prohibited bases apply to all programs). Persons with disabil audiotape, etc.) should contact the USDA's TARGET Center at 202-720-2600 (voice and TDD).	a me date necessa, and complet	and reviewing the collection of information.
To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 236 W. Mar.		